



# Haringey

## Local Tobacco Control Profiles

### Introduction

The aim of these profiles is to provide information for local government, health organisations, commissioners and other agencies to monitor the impact of tobacco on local communities, and to monitor the services and initiatives that have been put in place to tackle tobacco related problems.

The indicators contained within the web-tool were selected following consultation with stakeholders and a review of the availability of routine data. They are not exhaustive and should be used alongside local intelligence to gain a full picture of the range of tobacco control initiatives and issues in each locality.

The tobacco control profiles are part of a series of products produced by Public Health England that provide local data alongside national comparisons to support local health improvement.

This document is part of the Local Tobacco Control Tool. For further information about each indicator please view the definitions tab within [the tool](#). For further information on tobacco and health please [click here](#).

### Contents

#### Spine Charts

Key Indicators	Page 2
Smoking prevalence in adults	Page 2
Smoking prevalence in young people	Page 2
Smoking related mortality	Page 3
Smoking related ill health	Page 3
Impact of smoking	Page 4
Smoking quitters	Page 4

#### Tartan Rugs

Key Indicators	Page 5
Smoking prevalence in adults	Page 5
Smoking prevalence in young people	Page 6
Smoking related mortality	Page 6
Smoking related ill health	Page 7
Impact of smoking	Page 7
Smoking quitters	Page 8

#### Summary Charts

Key Indicators	Page 9
Smoking prevalence in adults	Page 11
Smoking prevalence in young people	Page 13
Smoking related mortality	Page 16
Smoking related ill health	Page 17
Impact of smoking	Page 19
Smoking quitters	Page 21

#### Trends

Key Indicators	Page 22
Smoking prevalence in adults	Page 23
Smoking prevalence in young people	Page 24
Smoking related mortality	Page 26
Smoking related ill health	Page 27
Impact of smoking	Page 28
Smoking quitters	Page 29

[www.tobaccoprofiles.info](http://www.tobaccoprofiles.info)


Produced by Public Health England.

For enquiries please contact [LondonKIT@phe.gov.uk](mailto:LondonKIT@phe.gov.uk) or call us on 020 7654 8172

# Spine Charts

## How to interpret the spine charts

- Significance compared to England average:
- Significantly worse
  - Significance not tested
  - Not significantly different
  - ◆ Regional average
  - Significantly better



## Key Indicators

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking Prevalence in adults - current smokers (IHS)	2014	20.7	18.0	26.9		9.8
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014	39.1	28.0	41.1		14.0
Successful quitters at 4 weeks	2014/15	1902	2829	957		5741
Smoking status at time of delivery	2014/15	- x	11.4	27.2		2.1
Smoking attributable mortality	2012 - 14	271.4	274.8	458.1		184.9
Smoking attributable hospital admissions	2014/15	1853	1671	2835		1030
Deprivation score (IMD 2010)	2010	36.1	21.7	5.4		43.4

## Smoking prevalence in adults

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking Prevalence in adults - current smokers (IHS)	2014	20.7	18.0	26.9		9.8
Smoking Prevalence in adults - ex-smokers (IHS)	2014	31.3	33.9	14.4		44.2
Smoking Prevalence in adults - never smoked (IHS)	2014	47.9	48.1	38.1		65.0
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014	39.1	28.0	41.1		14.0
Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS)	2014	15.1	30.8	12.7		40.9
Smoking prevalence in adults in routine and manual occupations - never smoked (IHS)	2014	45.8	41.2	29.1		61.9
Smoking prevalence in adults - current smokers (QOF)	2014/15	21.0	18.6	27.1		12.3
Smoking status at time of delivery	2014/15	- x	11.4	27.2		2.1

## Smoking prevalence in young people

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking prevalence age 15 years - regular smokers (SDD survey)	2014		8			
Smoking prevalence age 15 years - occasional smokers (SDD survey)	2014		5			
Smoking prevalence at age 15 - current smokers (WAY survey)	2014/15	6.5	8.2	14.9		3.4

Note: x - Value Missing

Smoking prevalence in young people continued	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking prevalence at age 15 - regular smokers (WAY survey)	2014/15	3.5	5.5	11.1		1.3
Smoking prevalence at age 15 - occasional smokers (WAY survey)	2014/15	3.1	2.7	7.6		0.6
Use of e-cigarettes at age 15 years (WAY survey)	2014/15	10.2	18.4	33.9		7.2
Use of other tobacco products at age 15 years (WAY survey)	2014/15	23.5	15.2	28.8		5.5
Smoking prevalence at ages 11-15 - regular smokers (modelled estimates)	2009 - 12	2.5	3.1	4.7		1.1
Smoking prevalence at ages 11-15 - occasional smokers (modelled estimates)	2009 - 12	1.0	1.4	2.0		0.5
Smoking prevalence at age 15 - regular smokers (modelled estimates)	2009 - 12	7.1	8.7	12.7		3.2
Smoking prevalence at age 15 - occasional smokers (modelled estimates)	2009 - 12	2.7	3.9	5.3		1.4
Smoking prevalence at ages 16-17 - regular smokers (modelled estimates)	2009 - 12	12.3	14.7	20.7		5.7
Smoking prevalence at ages 16-17 - occasional smokers (modelled estimates)	2009 - 12	4.1	5.8	7.8		2.2

## Smoking related mortality

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking attributable mortality	2012 - 14	271.4	274.8	458.1		184.9
Deaths from lung cancer	2012 - 14	54.2	59.5	107.7		29.8
Deaths from chronic obstructive pulmonary disease	2012 - 14	45.5	51.7	103.6		28.1
Smoking attributable deaths from heart disease	2012 - 14	30.8	29.7	58.1		18.9
Smoking attributable deaths from stroke	2012 - 14	12.4	9.3	17.8		6.3

## Smoking related ill health

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Premature births (less than 37 weeks gestation)	2010 - 12	80.3	75.8	107.9		60.6
Low birth weight of term babies	2014	3.1	2.9	5.8		1.6
Hospital admissions for asthma (under 19 years old)	2013/14	179.8	197.1	509.1		54.6
Smoking attributable hospital admissions	2014/15	1853	1671	2835		1030
Cost per capita of smoking attributable hospital admissions	2011/12	40.5	38.0	59.3		23.0
Lung cancer registrations	2010 - 12	70.0	76.0	146.8		40.1
Oral cancer registrations	2010 - 12	15.8	13.2	21.5		8.1

## Haringey

## Impact of smoking

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Indicative tobacco sales figures (£ millions)	2013	74.9	15446.1	440.2		13.2
Illicit tobacco: tax gap (£ millions)	2014/15		1200			
Illicit tobacco: share of cigarette sales (%)	2014/15		10			
Fatalities from accidental fires ignited by smoking materials	2013/14		72			
Accidental fires ignited by smoking related materials	2013/14		3300			
Attitudes to smoking in 15 year olds - 'smoking causes harm to others' (WAY survey)	2014/15	90.9	90.9	87.0		96.3

## Smoking quitters

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Number setting a quit date	2014/15	4279	5549	1955		12793
Successful quitters at 4 weeks	2014/15	1902	2829	957		5741
Successful quitters (CO validated) at 4 weeks	2014/15	969	1954	478		4953
Completeness of NS-SEC recording by Stop Smoking Services	2014/15	82.7	88.3	30.8		100
Cost per quitter	2014/15	340	420	1185		98

# Tartan Rugs

## Key Indicators

Period	England	London	Bark & Dag	Barnet	Bexley	Brent	Bromley	Camden	C London	Croydon	Ealing	Enfield	Greenwich	Hackney	Ham & Fulh	Haringey	Harrow	HaVering	Hillingdon	Hounslow	Islington	Kens & Chel	Kingston uT	Lambeth	Lewisham	Merton	Newham	Redbridge	Richmond	Southwark	Sutton	T Hamlets	Waltham For	Wandsworth	Westminster
Smoking Prevalence in adults - current smokers (IHS)	2014 18.0	17.0	21.7	13.2	16.6	13.6	14.0	16.9	13.3	17.1	16.4	13.6	17.3	20.0	22.2	20.7	13.1	18.4	17.1	12.3	22.2	18.7	13.5	18.1	20.6	15.5	20.6	14.2	11.2	16.5	15.3	22.1	20.7	14.4	20.0
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014 28.0	25.3	25.1	27.7	27.1	21.1	16.3	27.2	- x	18.9	26.5	17.8	29.4	23.9	33.2	39.1	20.7	27.2	30.9	21.6	41.1	26.7	15.1	23.4	19.5	20.2	27.5	17.3	14.0	23.4	25.5	36.8	23.7	36.7	28.7
Successful quitters at 4 weeks	2014/15 2829	3064	1587	1530	3040	2447	2453	3581	16884	4304	4914	4048	5228	2324	4693	1902	2313	3457	2810	5323	3159	2753	1720	2977	3303	2292	1937	3901	1486	2056	2548	3123	3591	3670	2109
Smoking status at time of delivery	2014/15 11.4 \$	4.8 \$	10.2	3.7	7.9	3.5	4.7	- x	- *	6.9	3.7	- x	- x	4.9 *	2.4	- x	4.7	10.4	7.4	3.3 \$	- x	2.2	3.9	3.4	4.9	4.4	5.0	- x	2.6	3.1	6.2	3.4	- x	3.8	2.1
Smoking attributable mortality	2012 - 14 274.8	261.4	365.0	202.9	269.7	219.9	237.2	246.7	- x	254.4	228.0	243.6	323.2	326.0	314.9	271.4	184.9	289.4	262.6	267.6	639.5	232.4	233.1	305.8	312.6	254.3	271.9	202.0	224.6	316.8	285.3	350.7	267.7	273.5	220.0
Smoking attributable hospital admissions	2014/15 1671	1517	1870	1243	1435	1333	1287	1536	878	1414	1621	1666	1650	1773	1871	1853	1239	1390	1610	1490	2306	1326	1373	1764	1968	1372	1496	1180	1186	1782	1387	1965	1380	1670	1294
Deprivation score (IMD 2010)	2010 21.7		34.2	16.6	16.7	30.5	15.0	25.4	11.1	22.8	25.0	26.1	31.9	42.9	27.5	36.1	15.5	16.6	19.8	21.8	35.9	23.3	11.7	31.2	31.0	14.6	41.8	20.4	10.1	29.7	15.4	39.6	35.4	21.5	24.6

## Smoking prevalence in adults

Smoking Prevalence in adults - current smokers (IHS)	2014 18.0	17.0	21.7	13.2	16.6	13.6	14.0	16.9	13.3	17.1	16.4	13.6	17.3	20.0	22.2	20.7	13.1	18.4	17.1	12.3	22.2	18.7	13.5	18.1	20.6	15.5	20.6	14.2	11.2	16.5	15.3	22.1	20.7	14.4	20.0
Smoking Prevalence in adults - ex-smokers (IHS)	2014 33.9	29.4	23.5	29.6	35.8	22.4	35.6	32.1	35.6	32.8	30.9	29.4	31.6	30.4	35.4	31.3	24.9	34.3	23.9	29.5	30.6	36.8	32.9	31.4	27.8	27.9	14.4	25.1	36.6	32.1	37.0	21.3	21.7	32.7	28.1
Smoking Prevalence in adults - never smoked (IHS)	2014 48.1	53.6	54.8	57.1	47.5	64.0	50.4	51.1	51.1	50.1	52.7	57.0	51.1	49.6	42.3	47.9	62.0	47.3	59.0	58.2	47.2	44.5	53.6	50.5	51.6	56.6	65.0	60.8	52.2	51.4	47.7	56.6	57.7	52.9	51.9
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014 28.0	25.3	25.1	27.7	27.1	21.1	16.3	27.2	- x	18.9	26.5	17.8	29.4	23.9	33.2	39.1	20.7	27.2	30.9	21.6	41.1	26.7	15.1	23.4	19.5	20.2	27.5	17.3	14.0	23.4	25.5	36.8	23.7	36.7	28.7
Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS)	2014 30.8	24.6	23.7	19.6	30.2	16.9	39.5	27.2	- x	34.2	25.4	26.2	25.5	24.3	34.7	15.1	27.5	26.4	21.1	20.7	21.7	32.7	38.1	23.1	27.0	19.5	15.6	25.6	34.8	23.8	33.0	14.6	24.0	12.7	24.3
Smoking prevalence in adults in routine and manual occupations - never smoked (IHS)	2014 41.2	50.1	51.2	52.8	42.7	61.9	44.2	45.5	- x	46.8	48.0	56.0	45.1	51.8	32.1	45.8	51.8	46.4	48.0	57.7	37.2	40.6	46.9	53.5	53.5	60.2	57.0	57.1	51.2	52.8	41.5	48.6	52.4	50.6	47.0
Smoking prevalence in adults - current smokers (QOF)	2014/15 18.6	18.1	20.8	15.1	16.9	17.6	15.9	17.9	11.2	17.9	16.2	19.1	19.9	22.7	19.9	21.0	13.4	16.8	17.0	16.8	21.4	19.0	16.1	20.6	20.5	16.7	18.7	14.1	13.6	19.3	17.3	21.7	20.5	17.0	17.3

Comparison with respect to England value Better Similar Worse Not compared

Note: \* - Disclosure control applied, x - Value Missing, \$ - Data quality note



## Smoking related mortality

	Period	England	London	Bark & Dag	Barnet	Bexley	Brent	Bromley	Camden	C London	Croydon	Ealing	Enfield	Greenwich	Hackney	Ham & Fulh	Haringey	Harrow	Havering	Hillingdon	Hounslow	Islington	Kens & Chel	Kingston uT	Lambeth	Lewisham	Merton	Newham	Redbridge	Richmond	Southwark	Sutton	T Hamlets	Waltham For	Wandsworth	Westminster
Smoking attributable mortality	2012 - 14	274.8	261.4	365.0	202.9	269.7	219.9	237.2	246.7	- x	254.4	228.0	243.6	323.2	326.0	314.9	271.4	184.9	289.4	262.6	267.6	339.5	232.4	233.1	305.8	312.6	254.3	271.9	202.0	224.6	316.8	285.3	350.7	267.7	273.5	220.0
Deaths from lung cancer	2012 - 14	59.5	57.7	84.0	45.6	60.4	51.3	47.7	57.7	- x	53.7	50.9	59.7	70.6	74.6	70.6	54.2	43.7	57.6	60.7	62.6	79.9	52.3	45.1	67.0	68.9	53.9	67.2	44.7	46.7	72.2	59.0	82.3	55.7	57.7	50.7
Deaths from chronic obstructive pulmonary disease	2012 - 14	51.7	50.8	89.5	34.3	46.3	34.9	43.3	49.0	- x	53.0	40.4	40.7	73.8	60.9	58.8	45.5	29.2	64.6	51.4	52.6	61.5	42.1	45.5	62.8	66.1	47.3	70.2	35.5	42.5	71.9	58.9	79.7	54.0	50.9	38.3
Smoking attributable deaths from heart disease	2012 - 14	29.7	27.7	35.6	22.1	29.0	28.4	25.2	21.2	- x	28.6	27.8	26.8	30.5	36.0	33.5	30.8	22.4	28.0	26.0	26.2	34.8	18.9	25.5	30.2	30.5	25.9	31.8	24.9	21.9	30.1	26.3	35.2	30.8	33.4	22.2
Smoking attributable deaths from stroke	2012 - 14	9.3	9.4	- x	7.2	8.5	10.8	8.0	- x	- x	8.1	8.9	8.8	9.9	12.7	- x	12.4	- x	9.3	9.3	9.0	- x	- x	- x	11.1	10.7	- x	12.2	8.8	- x	- x	10.1	13.8	9.1	9.3	9.8

## Smoking related ill health

Premature births (less than 37 weeks gestation)	2010 - 12	75.8	78.2	78.3	70.9	77.4	76.9	74.2	77.0	- *	83.3	80.2	87.5	79.9	75.3 *	71.5	80.3	81.0	77.1	94.2	72.4	71.2	77.0	64.1	72.3	89.2	66.1	96.8	75.9	69.4	83.5	66.1	73.9	82.7	70.3	75.1
Low birth weight of term babies	2014	2.9	3.2	3.4	2.5	2.9	3.6	2.6	2.9	- *	3.4	3.6	2.7	2.9	2.7 *	3.2	3.1	4.3	2.5	3.0	2.9	3.5	2.5	2.9	2.8	3.1	2.7	4.4	3.9	2.1	2.9	2.1	5.0	4.0	2.3	3.1
Hospital admissions for asthma (under 19 years old)	2013/14	197.1	204.8	166.1	108.9	165.2	271.5	140.3	145.8	-	355.4	256.1	146.8	223.3	243.3	112.3	179.8	260.2	126.6	149.1	156.0	235.3	154.9	101.2	307.0	236.7	246.5	226.5	209.9	80.6	259.1	191.1	171.0	360.4	206.4	119.8
Smoking attributable hospital admissions	2014/15	1671	1517	1870	1243	1435	1333	1287	1536	878	1414	1621	1666	1650	1773	1871	1853	1239	1390	1610	1490	2306	1326	1373	1764	1968	1372	1496	1180	1186	1782	1387	1965	1380	1670	1294
Cost per capita of smoking attributable hospital admissions	2011/12	38.0	39.1	50.2	35.5	39.8	35.3	38.5	41.6	- x	42.2	41.9	41.1	38.5	42.6	40.9	40.5	31.4	42.6	40.5	37.5	56.7	35.3	36.5	38.3	40.9	33.0	37.3	33.6	33.4	36.1	37.3	50.9	40.1	35.2	36.1
Lung cancer registrations	2010 - 12	76.0	72.2	109.2	58.1	72.9	61.8	55.8	75.4	77.9	66.7	56.7	66.1	86.8	91.9	87.8	70.0	47.8	76.3	66.5	72.4	104.0	66.0	55.5	84.4	90.0	73.8	89.7	54.3	59.0	97.7	73.9	116.5	68.8	81.3	73.1
Oral cancer registrations	2010 - 12	13.2	13.5	9.2	12.4	10.4	15.8	12.2	16.3	- *	13.3	15.1	13.2	13.7	9.9 *	20.2	15.8	12.4	8.2	14.6	14.3	14.9	15.4	13.1	15.3	13.3	11.7	14.9	10.5	12.7	10.2	13.2	21.5	13.0	14.8	17.9

## Impact of smoking

Indicative tobacco sales figures (£ millions)	2013	15446.1	2243.3	62.5	84.0	65.5	81.6	78.2	72.3	2.8	94.3	77.1	74.7	65.8	81.5	61.6	74.9	47.5	71.1	70.4	52.5	75.9	45.1	43.5	99.3	89.7	43.5	88.1	57.6	33.6	97.3	46.8	80.9	72.1	82.7	68.7
Illicit tobacco: tax gap (£ millions)	2014/15	1200																																		
Illicit tobacco: share of cigarette sales (%)	2014/15	10																																		

Comparison with respect to England value

Better	Similar	Worse	Not compared
--------	---------	-------	--------------

Note: \* - Disclosure control applied, x - Value Missing







# Summary Charts

## Key Indicators


Smoking Prevalence in adults - current smokers (IHS) (2014)




Smoking prevalence in adults in routine and manual occupations - current smokers (IHS) (2014)




Successful quitters at 4 weeks (2014/15)



Smoking status at time of delivery (2014/15)



Smoking attributable mortality (2012 - 14)




Key  England value and confidence interval  ↑ Haringey  London region value and confidence interval  
 Significantly worse  Not significantly different  Significantly better  Significance not tested


# Haringey

## Key Indicators continued

Smoking attributable hospital admissions (2014/15)




Deprivation score (IMD 2010) (2010)




Key    England value and confidence interval    Haringey    London region value and confidence interval  
 Significantly worse    Not significantly different    Significantly better    Significance not tested

## Smoking prevalence in adults


Smoking Prevalence in adults - current smokers (IHS) (2014)




Smoking Prevalence in adults - ex-smokers (IHS) (2014)




Smoking Prevalence in adults - never smoked (IHS) (2014)



Smoking prevalence in adults in routine and manual occupations - current smokers (IHS) (2014)



Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS) (2014)




Key   
 England value and confidence interval   
 Significantly worse   
 Not significantly different   
 Significantly better   
 Significance not tested   
↑ Haringey   
 London region value and confidence interval


# Haringey

## Smoking prevalence in adults continued


Smoking prevalence in adults in routine and manual occupations - never smoked (IHS) (2014)



Smoking prevalence in adults - current smokers (QOF) (2014/15)



Smoking status at time of delivery (2014/15)



Key     England value and confidence interval    ↑ Haringey     London region value and confidence interval  
 Significantly worse     Not significantly different     Significantly better     Significance not tested

## Smoking prevalence in young people


Smoking prevalence age 15 years - regular smokers (SDD survey) (2014)

No data


Smoking prevalence age 15 years - occasional smokers (SDD survey) (2014)

No data


Smoking prevalence at age 15 - current smokers (WAY survey) (2014/15)



Smoking prevalence at age 15 - regular smokers (WAY survey) (2014/15)



Smoking prevalence at age 15 - occasional smokers (WAY survey) (2014/15)




Key   
 England value and confidence interval   
 Significantly worse   
 Not significantly different   
 Significantly better   
 Significance not tested   
↑ Haringey   
 London region value and confidence interval


# Haringey

## Smoking prevalence in young people continued


Use of e-cigarettes at age 15 years (WAY survey) (2014/15)




Use of other tobacco products at age 15 years (WAY survey) (2014/15)




Smoking prevalence at ages 11-15 - regular smokers (modelled estimates) (2009 - 12)



Smoking prevalence at ages 11-15 - occasional smokers (modelled estimates) (2009 - 12)




Smoking prevalence at age 15 - regular smokers (modelled estimates) (2009 - 12)




Key  England value and confidence interval  ↑ Haringey  London region value and confidence interval  
 Significantly worse  Not significantly different  Significantly better  Significance not tested

## Smoking prevalence in young people continued


Smoking prevalence at age 15 - occasional smokers (modelled estimates) (2009 - 12)



Smoking prevalence at ages 16-17 - regular smokers (modelled estimates) (2009 - 12)



Smoking prevalence at ages 16-17 - occasional smokers (modelled estimates) (2009 - 12)




Key    England value and confidence interval    ↑ Haringey    London region value and confidence interval  
 ■ Significantly worse    ■ Not significantly different    ■ Significantly better    □ Significance not tested


# Haringey

## Smoking related mortality


Smoking attributable mortality (2012 - 14)




Deaths from lung cancer (2012 - 14)




Deaths from chronic obstructive pulmonary disease (2012 - 14)



Smoking attributable deaths from heart disease (2012 - 14)



Smoking attributable deaths from stroke (2012 - 14)




Key  England value and confidence interval  London region value and confidence interval  Significantly worse  Not significantly different  Significantly better  Significance not tested




## Smoking related ill health


Premature births (less than 37 weeks gestation) (2010 - 12)




Low birth weight of term babies (2014)




Hospital admissions for asthma (under 19 years old) (2013/14)



Smoking attributable hospital admissions (2014/15)



Cost per capita of smoking attributable hospital admissions (2011/12)




Key   
 England value and confidence interval   
 Significantly worse   
 Not significantly different   
 Significantly better   
 Significance not tested   
↑ Haringey   
 London region value and confidence interval


# Haringey

## Smoking related ill health continued

Lung cancer registrations (2010 - 12)




Oral cancer registrations (2010 - 12)



Key    England value and confidence interval    Haringey    London region value and confidence interval  
 Significantly worse    Not significantly different    Significantly better    Significance not tested

## Impact of smoking

Indicative tobacco sales figures (£ millions) (2013)



Illicit tobacco: tax gap (£ millions) (2014/15)

No data

Illicit tobacco: share of cigarette sales (%) (2014/15)

No data

Fatalities from accidental fires ignited by smoking materials (2013/14)

No data

Accidental fires ignited by smoking related materials (2013/14)


No data

Key   
 England value and confidence interval   
 London region value and confidence interval   
 Significantly worse   
 Not significantly different   
 Significantly better   
 Significance not tested




## Smoking quitters


Number setting a quit date (2014/15)




Successful quitters at 4 weeks (2014/15)




Successful quitters (CO validated) at 4 weeks (2014/15)



Completeness of NS-SEC recording by Stop Smoking Services (2014/15)



Cost per quitter (2014/15)




Key   
 England value and confidence interval   
 Significantly worse   
 Not significantly different   
 Significantly better   
 Significance not tested   
↑ Haringey   
 London region value and confidence interval


# Trends

## Key Indicators


Smoking Prevalence in adults - current smokers (IHS) - %




Smoking prevalence in adults in routine and manual occupations - current smokers (IHS) - %




Successful quitters at 4 weeks - per 100,000 smokers aged 16+




Smoking status at time of delivery - %




Smoking attributable mortality - per 100,000



Smoking attributable hospital admissions - per 100,000




Deprivation score (IMD 2010)




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

## Smoking prevalence in adults


Smoking Prevalence in adults - current smokers (IHS) - %




Smoking Prevalence in adults - ex-smokers (IHS) - %




Smoking Prevalence in adults - never smoked (IHS) - %




Smoking prevalence in adults in routine and manual occupations - current smokers (IHS) - %




Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS) - %




Smoking prevalence in adults in routine and manual occupations - never smoked (IHS) - %



Smoking prevalence in adults - current smokers (QOF) - %



Smoking status at time of delivery - %




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested


# Haringey

## Smoking prevalence in young people


Smoking prevalence age 15 years - regular smokers (SDD survey) - %




Smoking prevalence age 15 years - occasional smokers (SDD survey) - %




Smoking prevalence at age 15 - current smokers (WAY survey) - %




Smoking prevalence at age 15 - regular smokers (WAY survey) - %




Smoking prevalence at age 15 - occasional smokers (WAY survey) - %




Use of e-cigarettes at age 15 years (WAY survey) - %




Use of other tobacco products at age 15 years (WAY survey) - %




Smoking prevalence at ages 11-15 - regular smokers (modelled estimates) - %




Smoking prevalence at ages 11-15 - occasional smokers (modelled estimates) - %




Smoking prevalence at age 15 - regular smokers (modelled estimates) - %



Smoking prevalence at age 15 - occasional smokers (modelled estimates) - %



Smoking prevalence at ages 16-17 - regular smokers (modelled estimates) - %




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested



## Smoking prevalence in young people continued

Smoking prevalence at ages 16-17 -  
occasional smokers (modelled estimates) - %




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested


# Haringey

## Smoking related mortality


Smoking attributable mortality - per 100,000




Deaths from lung cancer - per 100,000




Deaths from chronic obstructive pulmonary disease - per 100,000



Smoking attributable deaths from heart disease - per 100,000




Smoking attributable deaths from stroke - per 100,000




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

## Smoking related ill health


Premature births (less than 37 weeks gestation) - per 1000




Low birth weight of term babies - %




Hospital admissions for asthma (under 19 years old) - per 100,000




Smoking attributable hospital admissions - per 100,000




Cost per capita of smoking attributable hospital admissions - £ per capita



Lung cancer registrations - per 100,000



Oral cancer registrations - per 100,000




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

# Haringey

## Impact of smoking


Indicative tobacco sales figures (£ millions) - £



Illicit tobacco: tax gap (£ millions) - £

No data available

Illicit tobacco: share of cigarette sales (%) - %




Fatalities from accidental fires ignited by smoking materials

No data available

Accidental fires ignited by smoking related materials

No data available


Attitudes to smoking in 15 year olds - 'smoking causes harm to others' (WAY survey) - %




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

## Smoking quitters


Number setting a quit date - per 100,000 smokers aged 16+




Successful quitters at 4 weeks - per 100,000 smokers aged 16+




Successful quitters (CO validated) at 4 weeks - per 100,000 smokers aged 16+



Completeness of NS-SEC recording by Stop Smoking Services - %



Cost per quitter - £



Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested